



The role of hydrogen bonds in the melting points of sulfonate-based protic organic salts

Luo, Jiangshui

Published in:
ISPE-XV, Uppsala 15-19th August 2016

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Luo, J. (2016). The role of hydrogen bonds in the melting points of sulfonate-based protic organic salts. In *ISPE-XV, Uppsala 15-19th August 2016* (pp. 177-177). [P52]

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The role of hydrogen bonds in the melting points of sulfonate-based protic organic salts

Jiangshui Luo^{1,2,3*}

¹ Department of Materials Engineering, KU Leuven, Leuven 3001, Belgium

² Centre for Surface Chemistry and Catalysis, KU Leuven, Leuven 3001, Belgium

³ Department of Energy Conversion and Storage, Technical University of Denmark, Denmark

jiangshui.luo@mtm.kuleuven.be

There are three main types of interactions inside organic salts – electrostatic interaction, hydrogen bonding and van der Waals force [1–4]. While van der Waals force is relatively weak, it is hydrogen bonding and particularly electrostatic interaction that determine the lattice energies of ionic systems and other physicochemical properties like melting points [5]. The hydrogen bond is an attractive interaction between a hydrogen atom from a molecule or a molecular fragment X–H in which X is more electronegative than H, and an atom or a group of atoms in the same or a different molecule, in which there is evidence of bond formation [6]. Hydrogen bonds in the solid state fall into the classification of strong, moderate, and weak hydrogen bonds [7]. In *molecular systems* like H₂O (vs. H₂S) or NH₃ (vs. PH₃), strong hydrogen bonds lead to higher melting points. However, in organic salts, the situation may be different [8,9].

In this presentation, we will present our important discovery on the influence of hydrogen bonding on the melting points of sulfonate-based protic organic salts [9]. We found five pairs, among which we will discuss in detail results of two highly similar *protic organic salts* – 1,2,4-triazolium methanesulfonate and imidazolium methanesulfonate. Both of them are solid-state proton conductors. The electrostatic interaction energy was calculated based on the single crystal X-ray and neutron diffraction data, showing that 1,2,4-triazolium methanesulfonate has higher electrostatic interaction energy at various temperatures. Moreover, variable-temperature infrared spectra, deuterium isotope effects, and single crystal data indicate much stronger hydrogen bonding inside 1,2,4-triazolium methanesulfonate. Therefore, 1,2,4-triazolium methanesulfonate is expected to have a higher melting point than imidazolium methanesulfonate. However, surprisingly, 1,2,4-triazolium methanesulfonate exhibits a much lower melting point (134 °C) than imidazolium methanesulfonate (188 °C). Therefore, it is concluded that stronger hydrogen bonding inside 1,2,4-triazolium methanesulfonate results in a much lower melting point than imidazolium methanesulfonate.

References

- [1] T. L. Greaves and C. J. Drummond, *Chem. Rev.*, 2008, **108**, 206.
- [2] A. Matic and B. Scrosati, *MRS Bull.*, 2013, **38**, 533.
- [3] H. Weingärtner, *Angew. Chem., Int. Ed.*, 2008, **47**, 654.
- [4] R. Hayes, S. Imberti, G. G. Warr and R. Atkin, *Angew. Chem. Int. Ed.*, 2013, **52**, 4623.
- [5] L. Glasser, *Inorg. Chem.*, 2012, **51**, 2420.
- [6] E. Arunan, G. R. Desiraju, R. A. Klein, J. Sadlej, S. Scheiner, I. Alkorta, D. C. Clary, R. H. Crabtree, J. J. Dannenberg, P. Hobza, H. G. Kjaergaard, A. C. Legon, B. Mennucci, and D. J. Nesbitt, *Pure Appl. Chem.*, 2011, **83**, 1637.
- [7] T. Steiner, *Angew. Chem. Int. Ed.*, 2002, **41**, 48.
- [8] A. Wulf, K. Fumino and R. Ludwig, *Angew. Chem., Int. Ed.*, 2010, **49**, 449.
- [9] J. Luo *et al.*, manuscripts in preparation.